

R. JOHN ELLIS, JR., M.D.
LAWRENCE A. SCHAPER, M.D.
MARK G. SMITH, M.D.
G. JEFFREY POPHAM, M.D.
AKBAR NAWAB, M.D.
MICHAEL SALAMON, M.D.
MATTHEW R. PRICE, M.D.
DANIEL E. RUEFF, M.D.



ORTHOPAEDIC SURGERY
FRACTURES
JOINT REPLACEMENT
SPORTS MEDICINE

OSTEOCHONDRAL AUTOLOGOUS TRANSPLANT PROTOCOL (OATS PROCEDURE) FEMORAL CONDYLE

GENERAL GUIDELINES:

- CPM machine at 0°- 60° for the first few days, then increase by 5° increments per day as tolerated. Utilize CPM for at least 4 weeks unless otherwise instructed by M.D.
- Nonweightbearing to Toe Touch Weight Bearing with crutches with brace locked at 0° for 4-6 weeks (Specified by the M.D.).
- Regain full knee Extension as quickly as possible. Sleep in the brace locked at 0° until full extension is maintained.
- Brace locked in full extension for one week and then unlocked in 10-20° increments if full extension has returned and when adequate quadriceps control is established.
- Wean from brace at 6 weeks unless otherwise instructed by M.D.
- Elevate the entire leg. Do not place pillows under the knee for long periods of time.
- Modalities: PRN for pain and inflammation.
- Utilize cryotherapy for 20 to 30 minutes every hour for pain and swelling reduction.

WEEKS 0-4

- Restore patellar mobility.
- Restore knee extension (low load-long duration stretching).
- Hamstring/gastrocnemius stretches.
- Ankle ROM for swelling and DVT prevention.
- PROM of knee (Seated flexion, heel slides, wall slides). Goal of 90 degrees by week 3.
- 4-way open chain straight leg exercises.
- *Focus on knee remaining locked in concentric and eccentric phase. Increase reps and hold time initially, before adding resistance. No ankle weight until extensor lag is eliminated.*
- Modalities: PRN for pain and swelling.
- NMES for Quadriceps activation and re-education.

WEEKS 4-7

- Increase ROM as tolerated, to full range.
- Continue to increase hip resistance during open chain exercises as tolerated.
- May start light aquatic exercises once good Quadriceps control is established and wound is healed appropriately.
- Isometric hamstring strengthening on a ball.
- Initiate stationary bike for range of motion without resistance.
- Gait training and proprioceptive/balance exercises as weight bearing restrictions allow.

WEEKS 8-12

- Increase resistance with all exercises as tolerated.
- Increase leg strength allow for normal gait and walking longer distances.
- Start closed chain exercises as tolerated in a protected range. (1/3 Partial Squats, Leg press with light resistance and high repetitions, calf raises, proprioceptive exercises, BAPS board)
- Initiate Step up/down exercises without compensations.
- Increase resistance with stationary bike.
- Sport specific exercises and Sportsmetrics program.
- Initiate jogging program when allowed by M.D.
- Agility drills when tolerated.
- Increase Cardio as tolerated.