

R. JOHN ELLIS, JR., M.D.
MARK E. PETRIK, M.D.
LAWRENCE A. SCHAPER, M.D.
MARK G. SMITH, M.D.
G. JEFFREY POPHAM, M.D.
AKBAR NAWAB, M.D.
MICHAEL SALAMON, M.D.
MATTHEW PRICE, M.D.
DANIEL RUEFF, M.D.



ORTHOPAEDIC SURGERY
FRACTURES
JOINT REPLACEMENT
SPORTS MEDICINE

TOTAL HIP ARTHROPLASTY PROTOCOL

The following protocol should be used as a guideline for rehabilitation progression, but may need to be altered pending the nature and extent of the surgical procedure, healing restraints or patient tolerance.

- **Total hip precautions after surgery should be followed for 3 months and include:**
 - **Avoid hip flexion past 90°**
 - **Avoid internal rotation of the lower extremity**
 - **Avoid crossing the midline of the body**
 - **Avoid sitting on low, soft surfaces**
 - **Use a raised toilet seat for 6 weeks – 3 months (discussed with the surgeon at the 6 week visit)**
- Patient will be weight bearing as tolerated with a rolling walking or other piece of medical equipment as seen fit by the surgeon. Patient may progress from a walker to a straight cane when they can demonstrate equal weight distribution, adequate balance, and limited Trendelenburg gait or limp.
- Patients may feel uneven in their gait, this is typically due to an adductor spasm and resolves over time.
- Driving is prohibited for the first 6 weeks or until off pain medication and walking without a cane.
- TED hose should be worn during waking hours and removed at night time. They will need to be worn for 6 weeks post operatively
- Patients will have staples or Steri-strips over their surgical incision. Staples will be removed at 10-14 post-operatively, then Steri-strips will be applied for 7 days. When the Steri-strips are removed at approximately 21 days post-op, begin scar massage.
- The patient may shower with a waterproof bandage over the staples but a shower stool or tub bench is recommended for the first 3 months. No soaking in the bathtub.
- Do not use weight machines until at least 3 months after your surgery.
- Patient will attend physical therapy 2-3x/week for the first 6 weeks or until patient returns to the surgeon.

TOTAL HIP ARTHROPLASTY PROTOCOL

PHASE 1: INITIAL PHASE

Weeks 1-3

- Walking at home for about 5 minutes every hour
- Bike for approximately 10 minutes
- Quad sets
- Glut sets
- Heel slides
- Supine hip abduction progress to side lying
- Short arc quad
- Modified bridges
- Clamshells
- Large arc quad
- Calf raises
- Heel raises
- HS curls
- Mini squats
- Step ups – only if patient is pain free with weight bearing

PHASE 2: INTERMEDIATE PHASE

Weeks 4-6

Continue with previous or modified versions of previous exercises, but may add:

- Progress ambulation from straight cane to no assistive device when patient can walk without a Tredenburg gait/limp
- Side lying abduction
- Marching
- Standing 4 way hip with theraband resistance
- Forward step ups – increase step height
- Lateral step ups – increase step height
- Single leg stance
- Wall sits
- Sit to stand without UE assistance
- Single knee to chest (at 6 weeks)

PHASE 3: ADVANCED PHASE

Week 7 – 3 months

Continue with previous or modified versions of previous exercises, but may add:

- Increase leg raise weight up to 5#
- Sidestepping with theraband resistance
- Walking program – start with ¼ mile and gradually increase
- Upper body weight machines
- For cardiovascular fitness – elliptical, walking outside or on a track, aquatic exercise, cycle. NO treadmill walking due to compression on the new joint.

PHASE 4: FINAL PHASE

3 months +

Continue with previous or modified versions of previous exercises, but may add:

- Begin using lower extremity weight machines
- Recommended activities – elliptical, cycle, walking, aquatic exercise, low impact aerobics, yoga, Tai Chi, Theraball exercises
- NOT recommended – running/jogging, high impact aerobics, jumping rope or plyometrics